

10 February, 2020



ISSN 2643-4776 (print) / ISSN 2643-4806 (online)

The Taxonomic Report

# Designation of neotype of *Hemileuca maia* (Drury, 1773) and refinement of its type locality (Bombycoidea, Saturniidae, Hemileucinae)

Harry Pavulaan 606 Hunton Place NE Leesburg, VA. 20176 intlepsurvey@gmail.com

**ABSTRACT**. The present paper defines the original type locality of *Hemileuca maia* (Drury, 1773) as Edgewood, New York, in the Long Island Pine Barrens. It has been locally known as the õCoastal Pine Barrens Buckmothö and its range may be confined entirely to Long Island, though populations throughout southeastern coastal New England and offshore islands may be considered the same nominotypical taxon. In analyzing historical events, there are questions regarding the geographic location, and source and eventual disposition of the specimen illustrated by artist Moses Harris in Drury (1773). While there is ample evidence that Druryøs specimen of *maia* came from a Mr. Thomas James of Brooklyn, New York, it is not clear whether the illustrated specimen was actually obtained by, and in the possession of, Dr. John Fothergill at the time the specimen was illustrated. Druryøs illustrated specimen perhaps found its way to various collections and is believed to now be lost or unidentifiable. A neotype is designated from the Edgewood Scrub Oak Plains, near present day Brentwood, New York, to agree with a possible collecting location used by Thomas James.

Additional key words: Pitch Pine Barrens, Scrub Oak Plains, Thomas James.

ZooBank registration: http://zoobank.org/References/6A0BCEEB-F6E2-4FC6-BFB8-BD116194AFA0

# **INTRODUCTION**

Dru Drury (1725-1803) stood out among entomologists in 19<sup>th</sup> century England as having likely the largest private natural history collection of the time, containing an estimated 11,000 specimens. Druryøs livelihood was as a jeweler and silversmith, but he was better known as an entomologist. He served as president of the Society of Entomologists of London (1780-1782) and was a member of the Aurelian Society. Among his peers were familiar names such as Cramer, Donovan, Fabricius, Kirby, and Linnaeus among others.

Drury produced and published his major work -Illustrations of Natural Historyø in three volumes: Vol. 1 (1770), Vol. 2 (1773), and Vol. 3 (1782), based on specimens in his collection. What is now known as *Hemileuca maia* was described by Dru Drury in Vol. 2 (1773) as *Phal[aena] Bomb[yx] Maia* (Fig. 1). Interestingly, species names were not included with any of the original descriptions nor the drawings in this work, thus the only reference to the name õmaiaö is found in the õIndex to the Second Volume. Names of the Insects, according to the System of Linnaeusö. Drury explained his curious philosophy on names: õIt is not my present design to enter into the scientific part of the study, by arranging the insects according to any system now established; nor will the reader find that I have given a single name to any one here figured. This, indeed, must be the consequence of not following the system of any author, unless I had formed one of my own; for it is impossible I should give names to them, particularly trivial ones.ö

# P L A T E XXIV.

#### FIG. III. Expands full two inches and a quarter.

Upper-fide. — The Antennæ are black, and ftrongly pectinated or combed. — The Neck is afh colour. — The Thorax and Abdomen black, the extremity being orange. — All the Wings are fimilar to each other, and very thin. — The Superior are black, with a white bar croffing them from the anterior to the posterior edges; whereon is a femi eye placed near the former. — The Inferior Wings are alfo black, with a white bar, broader than that on the fuperior wings, croffing them from the anterior to the abdominal edges; having near the former a black triangular fpot thereon, answering to the femi eye on the other wings. Under-fide.—The Palpi and Tongue, if it has any, are not to be feen.—The Legs and Thorax are black.—The Thighs orange.—The Abdomen grey, having its fides fpotted with white; the extremity being orange, as on the upper-fide.—All the Wings are the fame colour as on the upperfide, but rather more diftinct. The thinnefs of the wings occasion the colours to be lefs diftinct and clear, than in most others of this kind. The edges of all the wings are even.

It was brought from New York, and is in the possefition of Dr. Fothergill. I have not feen it in any author.

**Fig. 1.** Original description of *Phalaena Bombyx Maia* (in Drury, 1773). [Ed. Note: Placement of text is reformatted here, but content is intact].

The plate (Fig. 2) in Drury (1773) was drawn and engraved by artist and entomologist Moses Harris, whoøs engraving work was in high demand in England. Drury set a high standard for accuracy in the drawings: õ[T]he same plan, of giving *just* and accurate figures, that was followed in the first volume, is continued in this. The utmost care and nicety has been observed, both in the outlines and engraving. Nothing is strained, or carried beyond the bounds nature has set; and whoever will compare the engravings with the originals, I flatter myself will allow, that nothing is borrowed from fancy, or any colour given to an insect that does not really exist in the subject intended to be represented.ö One must assume that care was taken to represent exacting detail to width of the bands, placement of spots, etc. in the plate of *maia*. Of interest is the accurate depiction of the secondaries showing as a shadow, as seen through the primaries where the wings overlap.



Fig. 2. Phalaena Bombyx Maia (in Drury, 1773)



Fig. 3. Saturnia Maia (in Westwood, 1837)

Thirty-four years after the passing of his friend Drury, J. O. Westwood decided to republish Drury¢ work. In his Vol. 1 (1837), Westwood explained that Drury¢ work was of õacknowledged valueö, but the scarcity of Drury¢ work appeared to be õalmost unknown to Continental Entomologistsö. Thus, as Westwood stated: õI have consented to undertake the charge of bringing it forth in a form more adapted to the present greatly advanced state of Entomology.ö Which meant, in part, applying current names to each species treatment. He went on to write: õ[T]he chief difficulty has arisen from the non-possession of the specimens which served for the original illustrationsí ö. Unfortunately, all of the specimens illustrated in Drury¢ work had been sold or otherwise disappeared into the world of private and institutional collections by the time his collection was sold at auction in 1805. Typical of collections of the 18<sup>th</sup> and 19<sup>th</sup> centuries, generally only one or two specimens represented a species in any collection (Calhoun, 2010), which were easily õlost.ö Westwood mentions in a footnote that the copper plates, drawn and engraved by Moses Harris for Drury¢ 1773 work, were used exclusively in the first two volumes of Westwood¢s 1837 õNew Editionö. In Fig. 3 (above), one can see some minor differences in the result, as is expected with hand-colored illustrations.

Westwood, in Vol. 2 (1837), edited the description of *Phalaena Bombyx Maia*, and in this case applied the species name õ*Saturnia maia*ö (Fig. 4), referring to Druryøs *Phalaena Bombyx Maia* as a synonym. Westwood dispensed with the French translations that accompanied each of Druryøs original descriptions, added life history information such as hosts, descriptions of immature stages, flight dates, distribution, and anecdotes referring in this case to the name of common use in America: õBuck-flyö. Westwoodøs description of the adult is without any doubt representative of *H. maia*:

## SATURNIA MAIA.

Plate XXIV. fig. 3.

- ORDER: Lepidopters. SECTION: Nocturns. FAMILY: Bombycidæ, Leach.
- GENUS. Saturnis, Schrank. Attacus, Germar. Phalæna (Attacus), Drury.
- SATURNIA MAIA. Alis rotundatis nigris ; fascià albà, maculà subocellari nigrà, ano rufescenti. (Expans. Alar. 2 unc. 3 lin.)
- SIN. Phalæna (Bombyx) Maia, Drury, App. vol. 2. Cramer, Ins. 2. tab. 98. fig. A.
  Bombyx Proserpina, Fabr. Ent. Syst. 111. 1. p. 419. No. 40. Gmel. Linn. S. N. 2407. 480. Abbot & Smith Ins. Georg. pl. 50. Oliv. Enc. Méth. 5. 37. 48. Pal. Bauv. Ins. d'Afr. et d'Amer. Lep. pl. 24. f. 2. 3.
- HABITAT . New York (Drury). Georgia (Abbot).

Upper Side. Antennæ black, and strongly pectinated. Neck ash-coloured. Thorax and abdomen black, the extremity being orange. Wings pellucid. The anterior being black, with a white bar crossing them from the anterior to the posterior edges; whereon is a semi-eye placed near the former. Posterior wings black, with a broader white bar crossing them from the anterior to the abdominal edges; having near the former a black triangular spot thereon.

Under Side. Palpi and tongue indistinct. Legs and thorax black. Thighs orange. Abdomen grey, having its sides spotted with white; the extremity orange. Wings coloured as on the upper side, but rather more distinct. The thinness of the wings occasions the colours to be less distinct and clear than in most others of this kind. Margins of the wings entire.

Fig. 4. Description of Saturnia Maia (in Westwood, 1837).

At this point, one needs to determine the source and eventual disposition of the one original specimen that was modeled by Moses Harris for the plates of both the Drury (1773) and the Westwood (1837) works. A pertinent passage was written by Whitehead (1978) as follows: õ[T]he modern specialist trying to trace type or figured specimens must explore such famous old museums as the Leverian or Bullockøs and he must probe the many small and ephemeral collections that time and again changed hands, at each step with some loss of material and of the information that accompanied it. A complex web of sales, loans, gifts and transfers must be unraveled if the specimen is to be located or its provenance authenticated. The task is time-consuming, and it is often held that such õhistorical taxonomyö is an old-fashioned and expensive luxury in modern systematic work.ö Nevertheless, such a task must be pursued in the case of *Phalaena maia*.

It must be noted that Dru Drury enlisted the help of associates throughout the world to supply him with specimens and report natural history observations. Enlisting the employ of ship captains and other travelers, Drury sent letters and printed instructions abroad, even supplying collecting equipment when necessary, and always insisting on perfect specimens. Curiously, Drury (1773) mentions in the OD of *maia*: õlt was brought from New York, and is in the possession of Dr. Fothergill.ö This indicates that the specimen was not part of Druryøs own collection, at least at the time of writing Vol. 2.

Among Druryø overseas associates was Thomas James of Brooklyn, New York (Calhoun, 2010). Drury corresponded with James beginning in 1763, and received specimens from him between 1764 and 1776. Drury paid James for insect specimens and instructed him with collection techniques and sent him equipment. Drury had also encouraged James to collect duplicates, which Drury would exchange with other collectors. Thus there is a possibility there were additional specimens of *maia* being distributed, but no documented record of such transactions exists. It is believed that James originally lived in Flatbush, Brooklyn (New York), though some letters from Drury were addressed to James in õGuanoes, Long Islandö. The best current translation of õGuanoesö is similar to Gowanus (Brooklyn), a neighborhood near present-day Flatbush. Later correspondence between Drury and James went through a Mr. Rapalje, who lived at Brookland Ferry in the same area (Calhoun, 2010). The specimen of *maia* that James would have sent to Drury was collected or delivered in 1766, presumably while James still lived in Brooklyn. It is not known whether James could have collected the specimen of *maia* in Brooklyn or further out on Long Island, where James eventually relocated in 1767 or 1768.

The Gowanus and Flatbush areas at that time were primarily open farmland associated with scattered hamlets. The largest settlement of the time on the Brooklyn side of New York Harbor was around Brookland Ferry, where farmers from Long Island would take their produce across the East River to New York (Manhattan). This landscape was captured in excruciating detail in the õPlan of the City of New York in North America: surveyed in 1766 and 1767ö by Bernard Ratzer, commonly referred to as the õRatzer Mapö (Miller, 2016). Much of Brooklyn (Kings County) was apparently in agricultural production. East of this agricultural region, south of the glacial moraine, was a vast prairie-like plain stretching from present-day Queens County to Hempstead in central Nassau County (Flint, 1896), where a tiny remnant of the õHempstead Plainsö remains today protected, but otherwise unrecognizable to most. Earlier records documented the unbroken Pine Barrens of central and eastern Long Island extending westward only to around Hempstead (central Nassau County). As the Long Island population of *H. maia* is known to be primarily associated with extensive growth of Scrub Oak (*Quercus ilicifolia*), the likelihood exists that James collected Druryøs specimen farther east, not in Brooklyn.

Cockerell (1922) cited Druryøs correspondence through two letters with James, one of which confirms Jamesø move in that time period. In a letter dated Aug. 1, 1768 Drury requested James to send more insects from Jamesø new home out on Long Island, some 40 miles away, and included payment for specimens. Interestingly, the distance from Gowanus (Brooklyn) to the Edgewood Pine Barrens/Oak Plains, just east of present-day Deer Park, N.Y., is approximately 40 miles. In the mid-1700øs this area

was characterized by vast expanses of Pitch Pine/Scrub Oak forest. I have observed *H. maia* commonly in the Edgewood area since the late 1960¢s, when I lived in Brentwood, and collected it many times. Much of this area has now been converted to industrial use, though a considerable remnant of the Edgewood Pine Barrens and associated Scrub Oak Plains is protected within the New York State Dept. of Environmental Conservation¢s Edgewood Oak Brush Plains Preserve, and there are scattered remnants adjacent to the Preserve which will likely be developed in coming years.

Westwood¢s first hint at the disposition of Drury¢s specimens was as such: õ[A]s appears from the Catalogue of his [Drury¢s] Insects, which I obtained at the sale of Mr. [Edward] Donovan¢s collections, to whom, as appears by a note, they were presented by Mr. Druryí Here it is quite evident that Drury had carefully noted down the localities of all the specimens of this insect which he possessedí ö Drury reportedly produced printed circulars for the public sale of his collection in London, in 1788, several years prior to his death on Jan. 15, 1804, which included not only insects, but also cabinets, books and the original copper plates. Drury¢s handwritten collection catalogue provides whatever data was available to the author (Fig. 5).

Lepidoplera Thatena 10 06 Madres Mr. Sheene 1772 39 00 2 Paranympha New York - Mr. James. 1765 Linn p. 042 619 122 New York - Mr. James. 1765 Und Mustr Hol 2. M.241 30 06

**Fig. 5.** Portion of Druryøs handwritten catalogue confirming the source of the *maia* specimen to be õMr. Jamesö. It was apparently obtained in 1766. Note Drury listed it under õ*Phalena*ö [sic].

The original copy of Druryøs handwritten collection catalogue, four notebooks in his hand and presumably prepared around 1790, was presented to Edward Donovan, who in turn used it to produce the catalogue of Druryøs collection for public auction (Donovan, 1805; Calhoun, 2010), of which Westwood obtained a copy. It should be noted that *P. maia* was NOT specifically listed among sale items in the sale catalogue (though possibly grouped with several unspecified multiple õ*Phalaena*ö entries which were purchased by John Francillon). Westwood reproduced part of Druryøs handwritten catalog in his 1837 work to list purchasers of Druryøs collection. The following names were among those listed: G. Humphrey, Donovan, General Davies, Macleay, Latham, and Kirby. John Francillon was listed among those purchasing the multiple õ*Phalaena*ö entries, which could have included Druryøs specimen of *maia*. However, when Francillonøs own collection was auctioned in 1817-1818, there was no record of *maia* (Calhoun, pers. corr.). Interestingly, Macleay was the recipient of many of Druryøs specimens either through the sale of Druryøs collection or from the auction of Francillonøs collection.

The original specimen illustrated by Moses Harris (Drury, 1773) is believed to be either lost or unrecognizable among worldwide lepidoptera collections. A problem now encountered is that data labels were rarely affixed to specimens during the 18<sup>th</sup> and 19<sup>th</sup> centuries, and data (if any) was recorded in separate logbooks or catalogs (Hancock, 2004; Calhoun, 2010). Often, any labels affixed to specimen pins or trays were simply cabinet labels produced to identify a species. This information would have been imprecisely forwarded or lost in the process of specimen exchanges, thus becoming unreliable.

What further complicates matters is the mention in the OD of *Phalaena Bombyx Maia*, where Drury (1773) writes: õlt was brought from New York, and is in the possession of Dr. Fothergillö. It is not clear from whom John Fothergill obtained the specimen mentioned in the OD, but he apparently had his own contacts in the New York area who likewise sent him specimens. Fothergill is known to have received specimens from his Quaker friends in the New World (Douglas, 2015). Drury always indicated in his written work, who owned the specimen (Brock, 1977). This brings to question: Did Thomas James NOT collect the specimen described and illustrated in Drury (1773)? Calhoun (pers. corr.) suggests that Drury might have obtained additional specimens of *maia* at a later date and incorrectly credited them to James when he prepared his catalog. Also William Hunter is known to have received many American specimens from his own network of acquaintances and exchanged specimens with other collectors, among them a medical student of his, William Wood, who sent specimens mainly from Philadelphia and Rhode Island (Jeanne Robinson, pers. comm.; Calhoun, 2010). There was apparently considerable exchange of material back and forth between Drury, Fabricius, Macleay and Hunter (Hancock, 2004; Hancock et. al., 2015), often making it difficult or impossible to keep track of specimen transfers. Fothergilløs collection was bequeathed in 1780 to William Hunter, whose own collections formed the basis of the Hunterian Museum, which opened in 1807. Interestingly, the two õco-typesö of *Bombyx proserpina* Fabricius (Figs. 7 and 8), now housed in the Hunterian Museum (University of Glasgow), could possibly have come from Fothergill and served as the model for Druryøs Phalaena Bombyx Maia. Jeanne Robinson (pers. comm.) informed me that the only specimen data available comes from Fabricius (1775) where Fabricius mentioned the specimens came from America. The current cabinet labels were prepared between 1783 and 1785 by Hunterøs nephew, Matthew H. Baillie (Hancock et. al., 2015).

The two Hunterian specimens (Figs. 7 and 8) are a much closer match to Druryøs illustrated figure. Could one of these have been the model for Harrisø illustration for Drury? Fabricius described *proserpina* from Hunterøs collection in 1775. Calhoun (pers. corr.) advised this occurred five years before Hunter ever received Fothergilløs material. It is possible that Fothergill gave the original specimen to Drury. Both õco-typesö appear to be males by their broad antennae, though the University of Glasgowøs online collection entries indicate the sex of each is õunknownö. Druryøs specimen was obviously a male. Calhoun (pers. corr.) questioned why Drury would list a specimen of *maia* coming from James, yet possibly described the species from a Fothergill specimen. A satisfactory answer to this may never be found.

Fabriciusø description of *Bombyx proserpina* follows below (Fig. 6). The type locality is indicated as õHabitat in Americaö, but interestingly he lists Hunter (or the Hunter collection) as the source:



Fig 6: Description of *Bombyx proserpina* Fabricius (1775).

A rough translation of Fabricius (1775): õStature as follows. Antennae very comb-like, black, with a dark head. Thorax black, and the front fascia pale. The abdomen with dark reddish anus. The wings of the same colors, chimney-black, with white curves in the middle, in which a black eye is shaped by white. Black legs, thighs reddish brown.ö Fabricius refers to the abdomen having a dark reddish anus, characteristic of the male. Hunterian Museum specimen 133603 appears to be missing the orange-red terminal hairs, while 133604 is missing the abdomen entirely. The two specimens are nearly identical in phenotypic characters. A problem with these Hunterian Museum specimens is without detailed locality information they could be *H. maia*, or perhaps what is now known as *H. lucina* (characterized by broader white bands and greater translucence,) or possibly *H. nevadenis*, or another *Hemileuca* species. The bands are unusually wide for Long Island *maia*.



**Fig. 7.** Co-type of *Bombyx proserpina* Fabricius. Hunterian Museum catalogue number GLAHM 133603. Image reproduction by permission, © The Hunterian, University of Glasgow.



**Fig. 8.** Co-type of *Bombyx proserpina* Fabricius. Hunterian Museum catalogue number GLAHM 133604. Image reproduction by permission, © The Hunterian, University of Glasgow.

A specimen housed in the collection of the Macleay Museum (University of Sydney) (Fig. 9) might have passed through the hands of several people, thus losing any possibility of knowing its true source. An accompanying data label only indicates õNew Yorkö, and no other information accompanies the specimen. Robert Blackburn (pers. corr.) informed me that George Masters, first curator of the Macleay Museum, wrote the specimenøs data label. Such labels were written to replace the original labels. The specimen has also been re-pinned, as the current insect pin is not of the +old styleø type used by Drury. Possibly purchased by Macleay from the Drury collection, this specimen does not match the pattern of the specimen illustrated in Drury (1773), though the white bands are characteristic of Long Island *maia*. However, it does lend to the belief that several specimens of *maia* made their rounds among the collectors of the time. Since they are of unknown origin, it is unwise to designate a lectotype from among specimens of uncertain provenance.



**Fig. 9.** Male *Hemileuca maia* specimen housed in the Macleay Museum, Sydney, Australia. Image: © Macleay Collections, University of Sydney.

Also curious is the fact that a second *H. maia* specimen in the Macleay Museum collection, identified as õ*Hemileuca proserpina*ö indicates õGeorgiaö as the location. This specimen was likely collected by John Abbot, but it could not have been examined by Fabricius for the original description of *proserpina*, as Abbot did not arrive in Georgia until 1776.

The only other taxon of interest is the original holotype of *H. maia* õab. *lintneri*ö (Packard, 1914). The original description reads as follows:

õIn one [ab. *lintneri*] captured at Albany, N.Y. (fig. 11, Pl. LXVII), photographed by the late Dr. Lintner, there is no white band on the fore wings; the ocellus is distinct, but on the hind wing a white band incloses [sic] the ocellus.ö

A poor photograph is illustrated in Packard (1914), showing this extreme male variant of a *H. maia* male that is entirely missing the white band on the forewing. This specimen may quite possibly represent an aberrant specimen of *H. lucina*. The disposition of the specimen is presently unknown.

#### **NEOTYPE DESIGNATION**

Since none of the known historic specimens (despite links to Drury and his peers) clearly satisfy the phenotypic characters evident in the illustrated specimen (Drury, 1773), a neotype is proposed here to objectively define nominotypical *Hemileuca maia* (*Phalaena Bombyx Maia*) with a typical male, representative of the Long Island Pine Barrens population. I select a specimen from the Edgewood Pine

Barrens (Fig. 10). The Hunterian Museum specimens of *proserpina*, which are closer matches to the specimen illustrated in Drury (1773), do not appear to be of the nominotypical phenotype of *maia*. A specimen from the Long Island Pine Barrens region would best represent the species from the immediate õNew Yorkö (city) region, which is likely the geographical origin of the specimen illustrated in Drury (1773). The Long Island population is essentially a Pine Barrens isolate. The next nearest populations of *maia* occur in southeastern New England. Other populations from around Albany, N.Y. and in the New Jersey Pine Barrens region differ phenotypically by having more opaque, darker wings (Pavulaan, in press).



Fig. 10. Proposed neotype of Phalaena Bombyx Maia Drury (1773).

The proposed neotype, a male (Fig. 10), bears a plain white printed label [*Hemileuca maia* / October 21, 2017 / Long Island Avenue / north of Deer Park train station / Edgewood, New York. Collected by Harry Pavulaan] and a red printed label [NEOTYPE / *Phalaena Bombyx Maia* / Drury (1773) / Designated by / Harry Pavulaan 2020]. The neotype is deposited in the McGuire Center for Lepidoptera & Biodiversity, Gainesville, Florida.

## **TYPE LOCALITY**

The location of the defined type locality of the neotype of *Hemileuca maia* is just outside the east edge of the Scrub Oak Plains which lie within the Edgewood Oak Brush Plains Preserve, between Deer Park and Brentwood, Long Island, New York (Fig. 11). Industrial growth has destroyed most of the native forest cover of the area in the image with only patches of habitat remaining in some areas outside of the Preserve. This site, in what little remains of the historic hamlet of Edgewood, is directly north of the Long Island Railroadøs Deer Park station along Long Island Avenue. The type locality site will very likely be developed shortly after publication of this paper, though there is a thriving population of Buck Moths in the Scrub Oak Plains within the Preserve. A portion of the Scrub Oak habitat, which is visible in the image below (Fig. 12), was once part of the continuous Long Island Pitch Pine Barrens and Scrub Oak Plains, as recently as the mid-1960øs.



**Fig. 11.** Location of the defined type locality of the neotype of *Hemileuca maia* beyond the east edge of the Scrub Oak Plains within the Edgewood Oak Brush Plains Preserve between Deer Park and Brentwood, Long Island, New York.



**Fig. 12.** Typical view of Scrub Oak Plains habitat within the Edgewood Oak Brush Plains Preserve, immediately west of the proposed type locality. This is typical habitat for *H. maia* on Long Island. The Scrub Oak (*Q. ilicifolia*)  $\tilde{o}$ canopy $\tilde{o}$  in the foreground is the primary component of the Scrub Oak Plains and is approximately 5 ft. in height. Pitch Pine (*Pinus rigida*) towers above the Oaks in the background. [Photo: 10/21/2017]

#### ACKNOWLEDGMENTS

Many thanks to John Calhoun for helpful correspondence, pertinent reprints, direction to locating much of the historical sources used to write this paper, and for scanned images of *P. maia* from an original copy of Drury (1773), as well as that of *Saturnia maia* from an original copy of Westwood (1837), and for his review of the historical aspects of this paper; David Wright for final review and helpful comments; Jeanne Robinson (Curator of Entomology, Hunterian Museum) for discussions pertaining to exchanges between collectors in the late 18<sup>th</sup> century; Graham Nisbet (Rights and Reproductions Officer, Hunterian Museum) for image reproduction permission and improved image of *proserpina*; Robert Blackburn (Curatorial Assistant of the Macleay Museum) for discussions pertaining to the source of the Macleay Museum specimen and image reproduction permission; Jon Pelham for taxonomic discussion.

### LITERATURE CITED

- Brock, C. H. 1977. Dru Druryøs Illustrations of Natural History and the type specimen of *Goliathus goliatus* Drury. Journal of the Society for the Bibliography of Natural History 8(3): 259-265.
- Calhoun, J. V. 2010. Affirmation of the name *Papilio hyllus* Cramer (Lycaenidae) for a Nearctic butterfly, with the designation of a neotype. J. Lep. Soc. 64(2): 79-90.
- Cockerell, T. D. A. 1922. Dru Drury, an eighteenth century entomologist. Scientific Monthly 14(1): 67-82.
- Dononvan, E. 1805. A Catalogue of the Most Capital Assemblage of Insects Probably Ever Offered to Public Sale. Privately printed, London, U.K.: 16 pp.
- Douglas, S. 2015. Dr. John Fothergill: significant donor. Pp. 165-175. *In* Hancock, E. G., N. Pearce, & M. Cambell (eds.), William Hunterøs world: the art and science of eighteenth-century collecting. Ashgate Publishing Ltd, Surrey, UK.
- Drury, D. 1770. Illustrations of Natural History, Vol. 1. Printed by author, London: xxvii + 132 pp. + 51 pls, + 4 figs..
- Drury, D. 1773. Illustrations of Natural History, Vol. 2. Printed by author, London: vii + 94 pp. + 50 pls.
- Drury, D. 1782. Illustrations of Natural History, Vol. 3. Printed by author, London: xxviii +78 pp. + 50 pls.
- Fabricius, J. C. 1775. Systema Entomologiae, Sistens Insectorum Classes, Ordines, Genera, Species, Adjectis Synonymis, Locis, Descriptionibus, Observationibus. Flensburgh and Leipzig; Library Kotii, 8vo: xvi + 832 pp.
- Flint, M. B. 1896. Early Long Island: a Colonial Study. G. P. Putnamøs Sons, New York, N.Y.: 549 pp.
- Hancock, E. G. (2004). William Hunterøs insect collection and emerging descriptive taxonomy in the Eighteenth century. NatSCA News 4: 8-13.
- Hancock, E. G., N. Pearce, & M. Cambell (eds.). 2015. William Hunterøs world: the art and science of eighteenth-century collecting. Ashgate Publishing Ltd, Surrey, UK. 424 pp.
- Miller, L. 2016. Colonial New York Close Up: Revisiting Bernard Ratzerøs Plan of the City of New York. Brooklyn Historical Society. <u>https://www.brooklynhistory.org</u> (accessed 24 Jan. 2020).
- Packard, A. S. (*in*: Cockerell, T. D. A.). 1914. Monograph of the Bombycine Moths of North America, including their transformations and origin of the larval markings and armature (Part III) Families Ceratocampidae (exclusive of Ceratocampinae, Saturniidae, Hemileucidae, and Brahmaeidae. Memoirs of the National Academy of Sciences 12(1): ix + 516 pp.
- Westwood, J. O. 1837. Illustrations of Exotic Entomology, Containing Upwards of Six Hundred and Fifty Figures and Descriptions of Foreign Insects, Interspersed with Remarks and Reflections on Their Nature and Properties. By Dru Drury ó A New Edition, Brought Down to the Present State of the Science, with the Systematic Characters of Each Species, Synonyms, Indexes and Other Additional Matter. Henry G. Bohn, London. Vol. 1: xxvi + 123 pp. + 20 pl.; Vol. 2: vi + 100 pp. + 20 pl.; Vol. 3: vi + 94 pp. + 20 pl.
- Whitehead, P. J. P. 1978. A guide to the dispersal of zoological material from Captain Cookøs Voyages. Pacific Studies 2(1): 52-93.

# The Taxonomic Report is a publication of The International Lepidoptera Survey (TILS)

The International Lepidoptera Survey is registered as a non-profit Limited Liability Company (LLC) in the state of Virginia, U.S.A. The Taxonomic Report (TTR) is published for the purpose of providing a public and permanent scientific record. It appears in digital, open-access form, is regularly disseminated in hardcopy form to select institutional repositories and is also available as printed copy upon request at the discretion of authors and/or the editor. Printing and postage costs may apply. Contents are peerreviewed but not necessarily through the anonymous review and comment process preferred by some publishers of serial literature. An initial run of 25 copies is printed on paper to meet ICZN recommendation 8B. Copies of all TTR papers are available via the following digital repositories: Internet Archive (https://archive.org/). Biodiversity Heritage Library (https://www.biodiversitylibrary.org), Zobodat (https://www.zobodat.at/) and at the archival TTR website: (http://lepsurvey.carolinanature.com/report.html).

# **TILS Purpose**

TILS is devoted to the worldwide collection of Lepidoptera for the purpose of scientific discovery, determination, and documentation, without which there can be no preservation.

# **TILS Motto**

õAs a world community, we cannot protect that which we do not knowö

# Articles for publication are sought

They may deal with any area of research on Lepidoptera, including faunal surveys, conservation topics, methods, etc. Taxonomic papers are especially welcome. There are no page charges for authors. Before sending a manuscript, simply write to **TTR editor**, **Harry Pavulaan**, **606 Hunton Place NE**, **Leesburg**, **VA**, **20176**, **USA** to initiate discussion on how to best handle your material for publication, and to discuss peer review options; or email to <u>intlepsurvey@gmail.com</u>.

Visit *The International Lepidoptera Survey* on the World Wide Web at: http://lepsurvey.carolinanature.com

&

Join the discussion at our list serves on Yahoo! Groups at: http://groups.yahoo.com/group/TILS-leps-talk/ http://groups.yahoo.com/group/TILS-moth-rah/